

*Butorides
virescens*

FRENCH:
Heron vert
SPANISH:
Garcita verde

Green Heron

A relatively small, dark, compact, crested wading bird, the Green Heron is a common species of wetland thickets throughout much of North America. Although shy and retiring, it is a familiar sight to those spending time out of doors. Careful observers can see it stalking slowly through the water, perched quietly atop a branch, or as a dark form flying with slow wingbeats through the gathering dusk. Its flight call, an assertive *skeow*, is a sound typifying temperate and tropical wetlands of the Americas. Some of this bird's behaviors are especially well appreciated; flying away from human disturbance, for example, it often produces a scolding squawk and a stream of white defecation, giving it such vernacular monikers as "fly-up-the-creek," "shite-polk," and "chalk-line."

Although Green Herons generally nest solitarily, they sometimes nest socially in loose colonies. They feed day or night, standing patiently in shallow water waiting for fish, slowly stalking them through the shallows or diving on them from above. They are among the few tool-using birds, fabricating various baits that entice fish to where they can grab them.

Life Histories for
the 21st Century

The Birds of North America

Species limits within this small genus have been under intermittent review and revision, causing corresponding revisions in the "acceptable" English common name. The North American form of this species is part of a matrix of related populations, which differ





Figure 1.

Distribution of the Green Heron in North and Middle America. This species winters in small numbers north to Vancouver Island on the Pacific Coast. Closely related forms/species also breed in South America, Asia, Africa, and the Middle East. See text for details.

somewhat in size and plumage coloration, allocated to numerous subspecies but in total found over much of the tropical and subtropical world (Hancock and Kushlan 1984). At times similarities and purported intergradation of features among populations have been emphasized, leading to the North American form being lumped with Old World and Neotropical forms into a single species, *Butorides striatus*, most recently called the Green-backed Heron, more traditionally called the Striated Heron, and locally called the Mangrove Heron. At other times, differences and purported lack of intergradation of features among populations have been emphasized, leading to the North American form being separated from the Neotropical and Old World forms as *Butorides virescens*, called the Green Heron or Little Green Heron. On the Galapagos Islands, a third, distinctive population is sometimes considered a separate species, *Butorides sundevalli*, called the Galapagos or Lava Heron, but more often this population is viewed as a well-marked subspecies of the Neotropical form.

As this monograph is about the North American population, we call our bird the Green Heron (*Butorides virescens*), which conforms to the most recent declaration of the American Ornithologists' Union's check-list (Am. Ornithol. Union 1993). Because there is no evidence of fundamental differences in the biology (or indeed morphology) of any of the populations studied so far, however, we draw on information from *Butorides* herons worldwide.

DISTINGUISHING CHARACTERISTICS

Small, stocky heron (41–46 cm long), neck and legs short relative to elongated structures of many other herons. Adult plumages similar, but females tend to be smaller, duller, and lighter. Sexual dimorphism appears geographically variable according to older studies (Oberholser 1912). Adults with glossy greenish-black cap and back, wings black with greenish or bluish cast, underparts gray. Wing feathers edged in buff. Neck of adult is rufous—the character that distinguishes North American populations from those of *Butorides striatus*. Legs orange, bill dark. Shaggy greenish-black crest erectile. Young birds striped brownish on neck and below; back brownish with buff spots; shape and size distinguish them from young of any other species. Color pattern of both adults and young cryptic, clearly adaptive for undisturbed, solitary life in dense vegetation (Hancock and Kushlan 1984).

DISTRIBUTION

THE AMERICAS

Breeding range. Figure 1. E. North America from se. Canada (New Brunswick, s. Ontario and Quebec) south to Florida and Bahama Is.; west to e. North Dakota through Great Plains, avoiding drier areas of West, through s. Texas, Arizona, and New Mexico; along Pacific Coast from British Columbia through Baja California, south through mainland Mexico to s. Panama, Greater Antilles, Tobago, and islands off n. Venezuelan coast (Meyerriecks 1962, Peterson 1980, 1990, Hancock and Kushlan 1984). Range limited by high latitudes, aridity, and altitude. Local distribution constrained by suitable wetland marine or freshwater habitat for feeding.

Winter range. Figure 1. Postbreeding birds often disperse to areas where breeding does not occur. Most North American populations migratory; after postbreeding dispersal, most individuals from e. U.S. migrate south to winter along Gulf of Mexico, Florida, Caribbean islands, Mexico, through Central America to n. South America from ne. Colombia, Bonaire, and coastal Venezuela, infrequently as far as Guyana and Suriname. Small portion of northern populations winter north of Gulf Coast (e.g., Avakian 1984). Populations nesting in Florida and in southern states appear to be resident, but more data needed. Pacific Coast birds migrate southward to Mexico and Central America to El Salvador and Costa Rica, but many sedentary, some wintering as far north as Washington State.

OUTSIDE THE AMERICAS

Generally confined to AOU check-list area. Vagrants found in winter in Hawaii (Paton and MacIvor 1983). *Butorides striatus* populations widely distributed in tropical South America, sub-Saharan Africa, n. Madagascar, Indian Ocean islands, India, China, Japan, Southeast Asia through New Guinea, Australia, Pacific Ocean islands (Cramp 1977, Hancock and Kushlan 1984, del Hoyo et al. 1992).

HISTORICAL CHANGES

Sparse census data from early twentieth century obscure any long-term changes in distribution, but land development, drainage of wetlands, and recreational use of coastal wetlands may have reduced populations locally. Range expanding in midcontinent of North America along rivers and especially along impoundments, where species may be more common than reported because of solitary nature and cryptic plumage (Ryder 1979). Both breeding and

wintering range undergoing long-term expansion northward on Pacific Coast (Jewett 1939, Larrison 1940, 1947, Slipp 1942, Miller 1945, Eddy 1952, Peterson 1990).

FOSSIL HISTORY

Closely related *Butorides validipes* from early Pleistocene of Florida (Campbell 1976). But *B. virescens* also from Pleistocene of Florida (Wetmore 1931) and California (Howard 1936). Also from Native American kitchen middens about 500 yr old in California (DeMay 1942). Not clear that *B. virescens* has been distinguished or is separable from *B. striatus* in paleontological material. Amadon (1953) speculated that *B. virescens* may have reached North America from Asia, whereas *B. striatus* may have reached South America from Africa. But given related Pleistocene fossils and dispersal ability of these herons, historic distribution and dispersion patterns remain unclear.

SYSTEMATICS

GEOGRAPHIC VARIATION

Considerable geographic variation in plumage and size. Oberholser (1912) provided thorough documentation of the confusing level of plumage and size variation observed in museum specimens.

SUBSPECIES; RELATED SPECIES

Oberholser (1912) recognized 18 subspecies, mostly island forms. Most of these, including *patens* in area of *B. virescens*-*B. striatus* overlap in Panama, no longer recognized. Four subspecies currently accepted (Bent 1926, Payne 1974, Hancock and Kushlan 1984): *B. v. virescens*, central and e. North America to Panama; *B. v. anthonyi*, w. North America to n. Baja California, larger and paler; *B. v. frazari*, s. Baja California, with very dark neck; *B. v. bahamensis*, Bahama I., smaller and paler.

Relationships with related species a taxonomic yo-yo. Related to *Ardea*, *Egretta*, and *Nycticorax* herons (Sheldon 1987). At genus level, *Butorides* usually considered distinctive, confirmed by recent biochemical analysis (Sheldon 1987). Supposition that it is congeneric with *Ardeola* (e.g., Peters 1979) no longer supported.

Specific limits within genus are subject of considerable, ongoing debate, which we consider unresolved. *B. virescens* long considered a separate species (Wetmore 1965: 83-88) and member of superspecies with *B. striatus* and *B. sundevalli* (Bock 1956, Meyerrieks 1962). In 1970s

combined with *B. striatus*, based on morphological and behavioral similarities and analyses of individuals in area of overlap (Bock 1956, Curry-Lindahl 1971, Payne 1974, Am. Ornithol. Union 1976, 1983, Debenedictis 1978). More recently argued, based on same specimens, to be separate species (Sibley and Monroe 1990, Monroe and Browning 1992, Am. Ornithol. Union 1993, Debenedictis 1994). Debate probably not over. Additional studies of morphology and biochemistry still needed.

MIGRATION

NATURE OF MIGRATION IN THE SPECIES

Green Herons tend to wander after breeding, probably to seasonally more favorable foraging areas (Kushlan 1981). Dispersal not as extensive as in some heron species (Meyerrieks 1962) but may be intermittently eruptive (Hancock and Kushlan 1984), with stragglers as far as England (Cramp 1977). Dispersal merges gradually into protracted fall migration for northern birds. Gradual and variable character of dispersal/migration can be seen in records of summer fledglings banded in Maryland and Virginia. During October, these birds found as far north as Wisconsin and as far south as Nicaragua and Puerto Rico. Southern U.S. and some Pacific Coast birds may be mostly sedentary.

Migrates north in late winter and early spring, usually but not always at night, often in flocks, sometimes in large numbers (Scott 1890, Griscom 1923, Bent 1926, Meyerrieks 1962).

TIMING AND ROUTES OF MIGRATION

Fall migration begins late Aug-early Sep; most birds leave by mid-Oct (Bent 1926). Exact migration routes poorly known, but banded eastern birds have been recovered in Florida, Greater Antilles, Cayman I.; midcontinent birds in Texas, Florida, Mexico, Central America, Greater Antilles; western birds in Mexico. More banding data needed to establish precise migration routes.

Begins to arrive in ne. U.S. and California in Mar-Apr; e.g., in Ohio by mid- to late Mar (Peterjohn and Zimmerman 1989); in New York, "rare before mid-April" (Bull 1974); in Missouri, "not readily encountered" until late Apr (Robbins and Easterla 1992); in Vermont by mid-Apr (Laughlin and Kibbe 1985); in nw. U.S. by late May (Graf 1946). Arrives earlier than larger herons, perhaps because crepuscular feeding habits give birds longer feeding day (Seibert 1951).

MIGRATORY BEHAVIOR

Not reported.

CONTROL AND PHYSIOLOGY

Not reported. Temperature controls food availability (i.e., ponds freeze) and thus ultimately influences timing of migration. Likely that day length initiates migratory activities, but no data available.

HABITAT

BREEDING RANGE

Typically a bird of swampy thickets; has been said there is "scarcely a stream, swamp, or shoreline where it may not be found, whether fresh or salt" (Meyerrieks 1962). Although much exaggerated, this assertion suggests species' flexibility.

Nests in forest and swamp patches; may nest in dry woods or orchards, but usually near water (see Breeding: nest site).

Feeds in swamps, riparian zones along creeks and streams, also marshes, human-made ditches, canals, ponds, lake edges, open floodplains (*B. striatus*, Brown et al. 1982), backwater oxbow ponds, sloughs and side channels (Kaiser and Fritzell 1984), salt marshes (Clarke et al. 1984), mangrove swamps, pastures (Bryant 1914), mudflats, ponds in parks, and harbors. Although clearly prefers thick vegetation throughout range, will feed in open when food is available. In salt marshes, tends to hug creek banks; avoids open flats frequented by longer-legged herons.

SPRING AND FALL MIGRATION

Not specifically reported, but needs wetlands or coastal habitat for foraging on migration.

WINTER RANGE

Often coastal; uses all available marine and freshwater habitat. Mangrove swamp heavily used in tropical areas.

FOOD HABITS

FEEDING

Main foods taken. Carnivorous, typically a fish-eating species, despite comments to contrary in literature. However, prey selection is broad, depending on availability, and includes all sorts of invertebrates.

Microhabitat for foraging. Usually feeds by wading at water's edge or in very shallow water, reported as < 5 cm deep in Florida and New

Jersey and 1–10 cm deep in New York (Recher and Recher 1980). In closely related Mangrove Heron (*B. striatus macrorhynchus*), percent of 67 min foraging was: 4% in water 0–0.1 cm deep, 8% in 0.11–4.9 cm, 55% in 5–9.9 cm, 33% in 15–19.9 cm (Recher et al. 1983).

Also perches on branches, herbaceous vegetation, or rocks, close to water, where it accesses deeper water. Also feeds in deeper water by plunging into water after prey.

Generally under cover of thick vegetation. Despite this predilection, will feed in open, e.g., on exposed mudflats, tidal channels, open marshes, and pond edges.

Food capture and consumption. Heavy bill for bird's size, so it can capture large prey such as frogs, but not large fish that struggle (Recher and Recher 1980). Cryptic in its feeding habitat; dark dorsal and cryptic ventral color patterns match predictions of evolutionary-decision tree model for predicting plumage color of a species based on its feeding behavior and habitat use (Kushlan 1978).

Feeds at any time; during day, crepuscularly, also at night (Kushlan 1978, Recher et al. 1983, Hancock and Kushlan 1984, McNeil et al. 1993). Most arrivals at New Jersey roost occurred after sunset, and birds left in morning about 30 min before sunrise, before there was enough light to register on photometer (Seibert 1951).

Prey captured with darting stroke of head and neck, often with body lunge. When striking from horizontal, bird darts head and neck forward and down (Meyerrieks 1960). Prey generally grasped with bill but may be speared.

Green Heron (with Great Blue Heron [*Ardea herodias*]) has fewest recorded feeding behaviors of North American day herons (Kushlan 1976). Of 36 heron feeding behaviors, Green Heron (including *B. striatus*) known to use 15: Standing, Baiting, Standing Flycatching, Head Swaying, Neck Swaying, Walking Slowly, Walking Quickly, Scanning, Feetfirst Diving, Foot Stirring, Foot Raking, Plunging, Diving, Jumping, and Swimming Feeding (Kushlan 1976, Kushlan 1978, Davis 1983, Recher et al. 1983). Feeding activity index (measure of how actively bird moves while feeding) lower than that of any North American day herons and lower than that of nocturnal Black-crowned Night-Heron (*Nycticorax nycticorax*) (Kushlan 1978).

Most common feeding technique is to stand in crouched posture, body horizontal, head and neck retracted, looking into water. Will stand still for long periods before changing feeding sites. Birds stay longer in a spot after successful attempt than after unsuccessful one (Betts and Betts 1977).



Figure 2. Green Herons use bait to attract fish within striking distance, as seen here. They are among the few tool-using birds. Drawing by J. Zickefoose.

Standing is interspersed with Slow Walking, as bird moves slowly and methodically in crouched posture walking in water or along vegetation. When stalking prey, may move feet slowly, keeping 1 foot raised for up to 30 s. Foraging Mangrove Herons in 67 min of observation averaged 14 s/min walking, 13 steps/min, 0.9 steps/s for walking birds, 1 m/min distance moved, 0.2 prey attacks/min, and capture success rate of 57% (Recher et al. 1983).

Does feed more actively on occasion. In experiments with captive, bird sometimes Walked Quickly, perhaps to disturb hiding prey. Bird grabbed or speared fish, and had greatest capture success in shallowest water (76%, 0–10 cm) and poorest success in deepest water (20%, 20–30 cm) (Kramer et al. 1983). When feeding actively, often flips tail and raises and lowers crest feathers (Meyerriecks 1962). Foot Raking and Foot Stirring known but rarely used (Hoyt 1961). Raking has been recorded for both immature and presumably adult birds (Meyerriecks 1966, 1971).

May dive from perches head first into deep water, becoming submerged (Barker 1901). Diving is usually from perch, e.g., branch, pier, or rock.

Green Herons bait for fish (Fig. 2) with variety of baits and lures, e.g., crusts of bread (Lovell 1958), mayflies (Keenan 1981), and feathers (by

an immature bird; Norris 1975). For *B. striatus* in Japan, Higuchi (1986, 1988) reported that they bait fish more frequently in open water with few suitable perches. Three of these Japanese birds used lure bait (twigs, leaves, berries) 80% of time, live bait (flies and other insects) 20%. One bird spent 84% of foraging time baiting for fish; other birds used feathers and plastic foam for bait. Success rate highest when live bait used; juvenile birds did poorly baiting. One bird dug earthworms from mud and used them as bait, and twice birds broke pieces of stick to make bait, an example of tool-making. Baiting behavior rare among other herons.

DIET

Major food items. Opportunistic forager with varied diet (Wheelock 1906, Meyerriecks 1962, Kushlan 1978, Niethammer and Kaiser 1983, Hancock and Kushlan 1984). Overall diet includes invertebrates such as leeches; earthworms; aquatic and terrestrial insects such as adult and larval dragonflies (Odonata), damselflies (Zygoptera), waterbugs (Belostomatidae), diving beetles (Dytiscidae), grasshoppers, crickets, katydids (Orthoptera); spiders; crayfish, crabs, prawns; snails; fresh- and saltwater fish; frogs, toads, tadpoles, and newts; snakes and lizards; and rodents. Fish constitute primary food; these




Figure 3. Size and type of prey eaten by Green Herons in Louisiana during late summer. From Niethammer and Kaiser (1983).

include topminnows (*Fundulus*), minnows (*Notropis*), sunfish (*Lepomis*), catfish (Ictaluridae), pickerel (*Esox*), carp (*Cyprinus*), perch (*Aplodinotus*), gobies (Gobiidae), shad (*Dorosoma*), silverside (*Menidia*), eels (*Anguilla*), and in urban areas and human-made ponds, goldfish (Brooks 1923); 1 goldfish reported to be 10 cm and weigh 16 g (Hawbecker 1949). Will exploit a superabundant resource, e.g., grasshoppers in California (Bryant 1914) and frogs in New York (Recher and Recher 1980).

Quantitative analysis. Diet variability and opportunistic foraging behavior illustrated by analyses that follow and Appendix. Baynard (1912) reported for 1 Florida Green Heron stomach: 6 small crayfish, 16 grasshoppers, 2 cutworms, and small frogs. Meyerriecks (1962) reported U.S. Biological Survey data on 255 stomachs: 39% noncommercial fish, 6% food fish, 21% crustaceans, 24% insects, 10% other. Recher and Recher (1980) reported mean weight (g) of prey and (g/min hunting) in: Florida 0.7, (0.2); New Jersey 0.9, (0.3); New York 6.9, (3.9) (included frogs up to 125 g). Niethammer and Kaiser (1983) reported on 27 Green Heron stomachs from Louisiana containing 193 prey items, 143 of which were weighed and measured (Appendix, Fig. 3). Major insect types were dragonflies (Libellulidae), pygmy grasshoppers (Tetrigidae), and mole crickets (Gryllidae).

FOOD SELECTION AND STORAGE

Not reported.

NUTRITION AND ENERGETICS

Junor (1972) raised 2 *B. striatus* chicks on fish diet and reported that food requirements at flying stage were approximately 16% of body weight/d.

METABOLISM AND TEMPERATURE REGULATION

Chicks (*B. striatus*) are apparently seldom left uncovered during first 9–10 d after hatching; thereafter adults visit nest in daytime only to

feed chicks (Hindwood 1933). This suggests chicks achieve homeothermy by day 9 or 10. Cowles (1930) reported that before sunrise chicks crouched low in nest and huddled together, presumably for warmth. Later in day they became less active and breathing became rapid, with bills open, presumably to facilitate evaporative cooling. On hot days adults may shade eggs with wings (Cooley 1942).

DRINKING, PELLET-CASTING, AND DEFECTION


Adults often defecate while flying, particularly after being disturbed (Gordon 1961). Hindwood (1933) reported that scarcely any attempt is made by chicks to avoid defecation in nest, although others report that older chicks defecate over side of nest. Defecating bird lowers posterior, elevates tail, head, and neck, and droops wings slightly (Meyerriecks 1960).

SOUNDS

VOCALIZATIONS

Development. From hatching until fledging, chicks utter *tik-tik-tik-tik* (Meyerriecks 1962) or *toc-toc-toc-toc* (Hindwood 1933). Because these calls are given most intensively when chicks have not been recently fed, and peak when adults arrive at nest, they are best considered food-begging calls.

Vocal array. Call variously rendered as *skow* or *skeow* (Hancock and Kushlan 1984), *peu-ah* (Bent 1926), or *skyow* (Meyerriecks 1960). *Skow* slurs down from B₅ to F₅ or C₆ to G₅ (Meyerriecks 1962). Given as flight call and also as alarm call, the 2 being variously reported because of similarities. See Figure 4. During attack by a hawk, a Green Heron gave repeated "squawks" (Becker and Byers 1976). When disturbed at nest, birds utter *ku-ku-ku-ku-ku-ku* (Meyerriecks 1962),



skuk-skuk, kuk-kuk, or strong, hostile *raah-raah* calls (Meyerriecks 1960). When frogs jumped into a stream, a Green Heron emitted "squawks" before attacking frogs (Kalter 1932).

During breeding season, advertising calls or song by male are *show-ch* or *ow-ch* (Townsend 1928). Low-pitched *skow* call given from top of tree or other perch with bill up at 45° angle. Female advertising call is similar to alarm call, *skeow* or *skyow*. Attack calls a series of *raahs* (Meyerriecks 1960).

Phenology. Calls year-round. Low temperatures, wind, and rain reduce advertising and courtship displays, and hence reduce vocal component (Meyerriecks 1960).

Daily pattern of vocalizing. No data reported.

Places of vocalizing. *Skow* calls usually given in flight. Advertising call given from exposed perch or old nest.

NONVOCAL SOUNDS

Bill-snapping (clicking mandibles together) by males before copulation. Interpreted as interaction of hostile and sexual elements (Meyerriecks 1960).

BEHAVIOR

LOCOMOTION

Walking, hopping, climbing. Slow and deliberate when foraging; head and neck retracted or extended, sometimes tarsus practically on ground. Walks slowly or quickly over branches and roots, hopping from 1 to another. When walking, often erects and depresses crest and flips tail up and down (Bent 1926). On ground, will often walk to shrub and jump to top before flying (Meyerriecks 1960).

Flight. Slow, steady wingbeat at 2.8 beats/s, or 3.8/s after being flushed (Blake 1948). Resembles a tail-less crow; crows, when climbing, have similar wingbeat (2.7 beats/s). Normally flies with head retracted and legs extended beyond body, but often makes short flights with head and neck extended. Flight speeds reported: 35 kph (Wood 1933), 54 kph (McLean 1930), 40 kph Meyerriecks (1960).

When taking off, feathers sleeked, legs bent, posture horizontal with wings slightly extended, head and neck partially retracted, tail flipped or fanned laterally; then body thrust forward and upward by extending legs, wings extended upward, tail depressed, head and neck extended (crest erect or sleeked); then legs and feet brought together and extended rearward, and finally head and neck retracted. When landing, may make short glide, then head and neck extended to elevated position, crest erected, followed by dangling of legs which are then thrust forward and down, tail spread, wings beating as in hovering; perch grasped, head and neck retracted, crest lowered, body upright (Meyerriecks 1960).

Swimming and diving. Will swim after diving for prey (Hindwood 1933 for *B. striatus*). Web between middle and outer toes likely aids in swimming. One half-grown bird when placed in water swam gracefully 20 m, sitting erect "like a little swan" (Bent 1926).

SELF-MAINTENANCE

Preening, head-scratching, stretching, bathing, anting. Meyerriecks (1960) described maintenance behavior of Green Heron, which can reach entire body surface, except top of head and neck, with bill. Typical preening sequence begins with nibbling of breast powder-down patch; then grooms feathers, grasping each feather at base with bill tip, then moving smoothly upward or with rapid flicking movements. May shake with wings extended halfway, leaning forward, all feathers erect. Rubs head against body, may groom some or all feather tracts, repeatedly nibbling power-down feathers. Grooms scapular

Figure 4.

The flight/alarm call of the Green Heron. Top: perched bird. Bottom: in flight. Prepared by the Borror Laboratory of Bioacoustics (BLB), Ohio State University. Top: BLB #1514; bottom: BLB #14053.

plumes, raises and grooms neck feathers, nibbles and preens primaries with wing closed or partially open. Also preens under spread wing. Scratches head with pectinated claw of middle toe, crest raised. May peck toes. Ends bout with bill rubbing oil gland. Wipes bill on branch, alternating side of bill wiped. Stretches by extending head and neck, body forward, standing on 1 leg and extending opposite leg and wing. Yawns, especially after long incubation bout.

Powder down functions to dry and oil feathers in conjunction with oil gland (Hindwood 1933 for *B. striatus*). Of 3 pairs of powder-down feathers, pair on upper breast are largest, extending from near neck more than 6 cm. Second pair on rump, smallest pair on abdomen.

Able to tolerate high degree of attack by mosquitoes without employing defensive behavior (Edman et al. 1984). Corresponds with stealthy foraging behavior, where concealment is all important. When needed, will flick head and blink eyes to repel mosquitoes.

Sleeping and roosting. Sleeps with neck retracted, wings drooped slightly, body feathers somewhat erect, eyes closed. May sleep while brooding or incubating, head resting on nest rim. On ground may sleep with 1 leg retracted (Meyerriecks 1960).

May roost at night alone or in association with conspecifics or other colonial waterbirds. In mixed-species roosts, tends to segregate to a place adjacent to main roost. Up to 100 birds counted at one New Jersey roost (Seibert 1951).

Sunbathing. When sunbathing, droops half-open wings, raises head and neck slightly, usually faces sun, and may open bill, suggesting thermoregulation (Meyerriecks 1960). Hauser (1957) calls this "voluntary" or "normal" sunbathing, which often accompanies preening and other maintenance activities.

Daily time budget. No data reported.

AGONISTIC BEHAVIOR

Physical interactions. On feeding grounds, will defend a food resource (Meyerriecks 1960). A bird baiting fish with bread several times drove away American Coots (*Fulica americana*) with "threatening strokes of his long bill" (Lovell 1958).

On breeding grounds, if 1 bird gives Full Forward threat display (see below, Communicative interactions) and other bird displays in return, "a vicious fight ensues" (Meyerriecks 1960: 29). During fights birds peck at each other's heads, flail with wings, stab from horizontal stance. Supplanting attacks, where attacking bird replaces other bird from perch, common during early nesting season (Meyerriecks

1960). May attack and drive crows (*Corvus* spp.) away from eggs and nest, and usually intimidates grackles (*Quiscalus* spp.).

Communicative interactions. When humans approach nest, birds call aggressively, with harsh croaks, elongating neck, fluffing feathers, and raising crest (Dickerman and Gavino T. 1969). In conflict situations, erect crest, often tail-flip. In Forward Display, heron faces opponent, body horizontal, head and neck retracted, crouched; crest, neck, breast, back, flank feathers, and scapular plumes moderately erect; bill horizontal and closed; slow tail-flipping. Full Forward Display like Forward, but head and neck fully forward, feathers fully erect, eyes bulging, bill open showing red lining of mouth, vigorous tail-flipping; then lunges at opponent with wings slightly extended, uttering *raaah* calls; may make short flight toward opponent. In rare stiff-necked Upright Display, bird stands up with neck and crest feathers erect; may combine attack, escape, and sexual tendencies. Pursuit Flights on breeding grounds are probably low-intensity attack behaviors (Meyerriecks 1960).

SPACING

Territoriality. Defends feeding territories from conspecifics. In breeding season, song posts used to advertise territory, with most intense displays from old nest. These song posts mark courtship territories. In colonies and nesting aggregations, area defended around nest shrinks during nesting until, in some cases, only nest and immediately surrounding area are defended. Probably defends larger territories when nesting solitarily, but data needed.

Individual distance. No information.

SEXUAL BEHAVIOR

Mating system. All evidence, including high dispersion of nest sites, indicates seasonal monogamy. Given high degree of extra-pair mating in other colonial wading birds, information from marked birds or from biochemical studies would be desirable.

Pair bond, courtship displays, and mate-guarding. Meyerriecks (1960, 1962) described typical courtship display sequence. Begins with Flying Around displays, which resemble normal flight but are oriented to breeding sites, and with *skow* advertising calls from song posts. Pursuit Flight and Circle Flight displays accompanied by *skowing*, Crooked Neck Flight displays with partially retraced kinked neck; these displays have hostile component. Flap Flight Display is highest intensity of flight displays, where male

lurches through air with exaggerated flapping producing *whom-whoom-whoom* sound, neck crooked, legs dangling, *skowing*, crest, neck, and scapular feathers erect, sometimes giving *roo-roo* calls before landing. Soft parts change color in breeding condition, with bills glossy black, lores blue-black, legs bright red-orange.

Display flights interspersed with nonaerial displays such as Snap Display where male on perch points body, head, and extended neck below horizontal until bill tip at or below level of feet, crest and neck feathers slightly erect, and snaps mandibles together producing *click*. Variant includes bowing and bobbing movements. In Stretch display, male points bill straight up, stretches neck, and bends neck backward until head almost touches back, during which interscapular plumes are erect and fanned; then sways neck and head from side to side; crest, breast, and flank feathers sleeked, eyes bulge, and iris may change from yellow to deep orange; emits *aroo-aroo* sound.

Male performs Stretch before female is permitted to enter nest. Female performs Stretch after male, but female Stretch less intense. Stretch is employed during nest construction and nest relief. After female first enters nest, male's *skowing* and flight and Snap displays cease.

Copulation. Only on territory, usually on nest platform, and mostly during nest-building stage. First copulation usually on same day as pair formation. Before first copulation, birds engage in mutual bill-snapping and feather nibbling, but those behaviors reduced thereafter. Engage in Twig Passing and Stretch displays, some mutual, during copulation period. Typically, female in nest gives Stretch with back to male, who then mounts squatting female; both flap wings to maintain balance. Copulation lasts about 10 s (5–12); several hours between copulations. Copulations cease after last egg laid (Meyerriecks 1960, 1962).

Duration and maintenance of pair bond. Most authorities assume seasonal (serial) monogamy, but evidence from marked birds needed.

Extra-pair copulations. None reported.

SOCIAL AND INTERSPECIFIC BEHAVIOR

May nest in colonies with conspecifics or other colonial birds, generally somewhat apart from other species. In San Blas, Mexico, Burger (1978) found in colony of 320 pairs: 16% Tricolored Herons (*Egretta tricolor*), 13% Great Egrets (*Egretta alba*), 28% Boat-billed Herons (*Cochlearius cochlearius*), and 23% Green Herons, as well as 7% Olivaceous Cormorants (*Phalacrocorax olivaceus*)

and 13% Anhingas (*Anhinga anhinga*). For an Alabama colony, Gaston and Johnson (1977) reported Green Herons nesting with Tricolored and Little Blue (*Egretta caerulea*) herons, Snowy (*Egretta thula*) and Great egrets, and Glossy Ibis (*Plegadis falcinellus*). Also nests near grackles, which may help warn herons of predators or intruders (Wheelock 1906). Reported as an uncommon secondary species associated with roosts of European Starlings (*Sturnus vulgaris*) and Common Grackles (*Quiscalus quiscula*) (Caccamise and Fischl 1985).

Green Herons, although typically solitary foragers, foraged in New Jersey salt-marsh habitat with wide variety of waders, waterfowl, shorebirds, gulls, terns, raptors, and passerines (Burger et al. 1982).

May respond to presence of predator by assuming "Bittern Stance" with bill held vertical. Can see with binocular vision across base of bill from this position. May be cryptic adaptation for reed-bed environment (Cowles 1930 for *B. striatus*). During Bittern Stance, neck stretched upward, feathers sleeked, legs straight, eyes bulged, body motionless, except head may turn to follow predator.

Reacts to warning calls of many birds, including at least 10 Ardeidae species, crows, blackbirds, and grackles, and assumes alert posture, Bittern Stance, or flies. After pair formation, aggressive toward all passerines, especially crows and grackles. Considered intermediate in sociality between solitary breeding *Botaurus* bitterns and highly colonial herons, e.g., *Nycticorax* (Meyerriecks 1960).

PREDATION

Predation on eggs may be high, particularly by snakes (Dickerman and Gavino T. 1969, Kaiser and Reid 1987). Snake predation accounted for 25% of known nest failure in 1 study (Keiser and Reid 1987). Meyerriecks (1960, 1962) reported predation on eggs by Common Grackles and Fish Crows (*Corvus ossifragus*), and McVaugh (1975) reported probable predation by Boat-tailed Grackles (*Quiscalus major*).

Raccoons (*Procyon lotor*) probably take young birds (Dickerman and Gavino T. 1969). Becker and Byers (1976) reported unsuccessful attack on a flying Green Heron by a Broad-winged Hawk (*Buteo platypterus*). Innately recognizes some potentially dangerous animals. Six naive, hand-reared Panama birds fled from dead sea snake (*Pelamis platurus*), backing away with crests erect, then flew around aviary and tried to escape (Caldwell and Rubinoff 1983).

BREEDING

PHENOLOGY

Pair formation and nest-building. Figure 5. Pair formation may begin during migration; in New York both sexes arrive at breeding grounds simultaneously and begin breeding activities at once (Meyerriecks 1960, 1962).

Because of wide latitudinal range of breeding, initiation dates differ geographically. In North America, generally later farther north. Egg dates for Florida as early as 29 Mar, Gulf Coast first wk Apr, Midwest second wk May, New England and New York 29 Apr to late June (Bent 1926, Meyerriecks 1960), s. Ontario 4 May to mid-Jul (Peck and James 1983). In Oregon, birds nest Apr–Jun (Bent 1926, Graf 1946, Griffiee 1954).

In tropics, nesting season related to temperature and rainfall. In Veracruz, Mexico, in permanent aquatic habitat, nesting begins with increase in temperature—in Feb or Mar—but in San Blas, Mexico, begins in Jul with rainy season and flooding of mangrove flats (Dickerman and Gavino T. 1969).

Second brood per season. See Demography and Populations: measures of breeding activity: clutch size. Clutches for second nest begin soon after first is lost or completed; 5–9 d after loss of first clutch and as early as 9 d after successful nesting (Dickerman and Gavino T. 1969).

NEST SITE

Selection process. Nests in various habitats providing secluded nest sites and wetland feeding habitat: swamps, marshes, lakes, ponds, human-made or human-regulated habitat such as storm-water control impoundments and retention basins, even in dry woods and orchards in farmlands if feeding sites available (Bent 1926, Adams et al. 1985). Often nests on islands or tree islands in marshes. Nest is usually on or over water but may be up to 0.8 km from standing water (Kaiser and Reid 1987).

Nest may be placed alone, in loose aggregations, or in colonies. Kaiser and Reid (1987) suggested that birds nest alone if they can defend a feeding territory, e.g., along a stream, but nest in loose aggregations in wetlands where territorial defense is more difficult.

Single-species aggregations or colonies are small to moderate size. Examples include 70, 41, 17, and 8 pairs in Jamaica Bay, NY (Meyerriecks 1960, 1962). Nest densities variable. Of 137 nests in San Blas, Mexico, 45 ≤ 1 m apart, 75 1–2 m apart, and 21 > 2 m apart, with 5 nests only 15, 15, 20, 20, and 40 cm apart (Dickerman and Gavino T. 1969).

In mixed-species colonies, often nests some-




Figure 5.

Annual cycle of breeding, migration, and molt of the Green Heron in the United States. Thick lines show peak activity, thin lines off-peak. See Breeding: phenology, for a discussion of variation in the annual cycle of this species.

what apart from other species; usually well outnumbered by other species, e.g., 12 pairs in a mixed-species colony of 804 breeding pairs on Chimon I., CT (Marra and Bull 1986).

Nest site is chosen through process of territorial contraction. Male displays in and defends relatively large area that typically includes eventual nest site. As site is selected by pair and construction begins, pair concentrates its attentions on increasingly restricted area around nest.

Microhabitat. Nests in protected sites within trees or bushes. May nest in both aquatic and terrestrial sites in same area (Kaiser and Reid 1987).

Nest placed from ground level to 10 m above ground, occasionally to 20 m (Wheelock 1906, Huey 1915, Bent 1926, Howsley 1936, Cooley 1942, Mousley 1945, Meyerriecks 1962, Steirly 1964, Douglass et al. 1965, Dickerman and Gavino T. 1969, McVaugh 1975, Sandilands 1977, Osborn and Custer 1978, Coffey 1981).

Nests in many species of plants, and nest height determined in part by height and thickness of

plant. In San Blas, Mexico, for example, nests in 3 species of mangroves (red mangrove [*Rhizophora mangle*], black mangrove [*Avicennia nitida*], and particularly white mangrove [*Laguncularia racemosa*]); average height of 181 nests 0.68 m above water (Dickerman and Gavino T. 1969). On Ships Shoal I., VA, in shrubs 6–12 m high (Osborn and Custer 1978); also in Virginia, in loblolly pine (*Pinus taeda*; Steirly 1964). In Tennessee, in buttonbush (*Cephalanthus occidentalis*), 1–2 m above ground. In Arkansas, in post oak (*Quercus stellata*), 4 m above ground (Coffey 1981); also in Tennessee, in Virginia pine (*Pinus virginiana*), and cedars at heights of 4, 7, and 8 m, 5 m apart (Douglass et al. 1965). In Minnesota, in willow (*Salix* sp.) 1 m above water (Cooley 1942).

Meyerriecks (1962) reported nests in New York from ground level to 5 m above ground, most commonly near water, with concealment variable. In Magnolia, MA, colony of 20–30 pairs in pitch pines (*Pinus rigida*), 7 m above ground; in Westport, MA, colony of 20 nests on salt-marsh island in cedars, sassafras, and hickories, 1–7 m above ground (Bent 1926). In N. Carolina, nests in black willows (*Salix nigra*) over water (McVaugh 1975). In California, a nest in willow 8 m high (Howsley 1936). In Ontario, once nested in Wood Duck (*Aix sponsa*) box (Sandilands 1977).

In Missouri, height of 77 nests averaged 10.5 m above ground (range 2.2–19.0), primarily in honey locust but also in buttonbush, black willow, box elder (*Acer negundo*), and short leaf pine (*Pinus echinata*) (Kaiser and Reid 1987).

Site characteristics. Branches large enough to support nest, generally at a juncture. Birds appear to select trees with branching over those without. Overhanging branches to conceal nest nearly essential. Nest often suspended over water.

NEST

Construction process. Male selects nest site. Old nests reused in some places, after considerable renovation (e.g., Dickerman and Gavino T. 1969). Also builds in old nests of Black-crowned Night-Herons and Snowy Egrets (Meyerriecks 1962, Nickell 1966). In other places, old nests not reused, but sticks from nearby old nests refashioned into new nests.

Male initially begins nest construction. After pairing, male gathers nest material for female who does most of construction. Twigs worked into nest structure by "tremble-showing." Male typically breaks off twigs to pass to female. In early stages, twigs may be repeatedly dropped and retrieved. After eggs laid, incubating bird does most of building and repairing (Meyerriecks 1960). Birds continue to add new sticks through-

out nesting period and will pirate sticks from other active or old nests.

Building time for nest not reported in North America, but reported as 3 d and 7–14 d in *B. striatus* (Hindwood 1933, Brown et al. 1982).

Structure and composition matter. Nest round to elongate. Structure variable, from so thin that eggs are visible from beneath to substantial; total thickness of 72 nests averaged 11 cm, but recorded up to 24 cm thick, which may be reused nests (Dickerman and Gavino T. 1969).

Sticks used are relatively thin but may be quite long, up to 25 cm. Hindwood (1933) found about 220 sticks, as "thick as a lead pencil" and up to 46 cm long, in average nest of *B. striatus*. May also use reed, but doesn't use any lining material.

Dimensions. Diameter (outside) about 20–30 cm; Brown et al. (1982) reported 30 cm; mean diameter of solid part of 119 nests was 23 cm (Dickerman and Gavino T. 1969). However, a nest in Quebec was boat-shaped and diameter measured 36 x 20 cm outside (Mousley 1945).

Nest cup shallow; depth of nest depression about 0–10 cm, averaging 4.5 cm for 145 nests (Dickerman and Gavino T. 1969).

Microclimate. No data available.

Maintenance or reuse of nests, alternate nests. Old nest is focus of territorial behavior, e.g., most intense threat displays usually made on nest platform (Meyerriecks 1960, 1962). May reuse old nest platforms from previous season and refurbish with sticks from other old nests. Fifty-three of 59, and 67 of 68 nests surviving over winter reused following season in Long I., NY, colonies. Also uses twigs from old nests for new nests. Nests partially destroyed can be fixed rapidly, even while young hang on to nearby branches. One pair built nest to 0.25 m thick as water level rose threatening it (Dickerman and Gavino T. 1969). For second clutch in same season, may reuse same nest or make another (Dickerman and Gavino T. 1969). More likely to relocate nest following failure and to reuse nest following successful nesting (Kaiser and Reid 1987).

Nonbreeding nests. Not known.

EGGS

Shape. Elliptical to subelliptical.

Size. Length x breadth (mm): San Blas, Mexico ($n = 232$) 37.5 (range 34.0–40.8) x 29.2 (27.0–31.4) (Dickerman and Gavino T. 1969); New York ($n = 74$) 38.8 (35.7–43.3) x 29.6 (27.0–34.8) (Meyerriecks 1962); California ($n = 17$) 39.0 (37.0–42.0) x 29.5 (28.2–30.0) (Huey 1926).

Mass. No data.

Color. Initially clear, pale green to bluish, becoming duller and chalkier during incubation

period. May become stained with mud from adults. Color range called Lichen Green to pale Niagra Green of Ridgway (1912) (Dickerman and Gavino T. 1969). Unusual clutches of Olive Buff reported (Dickerman and Gavino T. 1969).

Eggshell thickness. Mean of 0.183 mm ($n = 302$) pre-pesticide era from 25 states; in regional comparisons with 1947–1973 data, changes from -1.2% to +1.6% ($n = 124$), none significant (Ohlendorf et al. 1979). Mean of 0.169 mm (SD = 0.011, $n = 8$) from Tennessee Valley in 1980–1981 (Fleming et al. 1984). Eleven percent thinning in eggs at Yazoo National Wildlife Refuge, MS, compared to pre-DDT era eggs (White et al. 1988).

Clutch size. Generally 3–5 eggs.

Egg-laying. Mostly in morning and early afternoon (Meyerriecks 1962, Dickerman and Gavino T. 1969). Clutches laid in 4–6 d, depending on clutch size. Second egg laid 2–5 d after first, in > 60% of cases laid on second day (Dickerman and Gavino T. 1969). Third egg laid 1–2 d after second, most commonly 2 d. Meyerriecks (1962) found less variability in New York.

INCUBATION

Onset of broodiness and incubation in relation to laying. Incubation typically begins with second egg, although first-egg start indicated in New York. Routine of "1 on, 1 off" incubation begins with second egg, but until third egg laid, incubation is intermittent.

Incubation patches. No data reported.

Incubation period. Somewhat variable, generally 19–21 d from last egg. Dickerman and Gavino T. (1969) reported 19 d for 8 nests, 20 d for 12 nests, and 21 d for 11 nests. Meyerriecks (1962) reported incubation period of 19 d for 1 nest, 20 d for 18, and 21 d for 1. Burns (1917) reported 17 d, probably too short. Because of timing of start, however, eggs in same nest not incubated for same period, ranging from 19 to 23 d in single nest (Dickerman and Gavino T. 1969)

Parental behavior. Both parents incubate and attend nest equally after full clutch. After third egg laid, male incubates mostly in midday and middle of night, with female mostly early morning and late evening, but schedule variable (Meyerriecks 1960).

Billing and feather-nibbling at nest relief, sometimes silent Stretch display, sometimes by both birds, early in incubation; later little or no nest relief ceremony. Both birds usually use same approach to nest, but if different, each bird consistently uses its own route (Meyerriecks 1962).

Hardiness of eggs. Eggs will die if exposed to sun and heat. Parents stand and shade eggs in

direct sun (Dickerman and Gavino T. 1969). This behavior most important in exposed nests. Eggs and young can die from heat. Neglected eggs can be lost to predators.

HATCHING

Preliminary events and vocalizations. Hindwood (1933) described hearing cheeping from chick and tap of egg tooth inside "slightly broken" egg, and "soft croaking" from another chick in perforated egg (in *B. striatus*).

Shell-breaking and emergence. Hatching reported at all daytime hours (Cooley 1942, Douglass et al. 1965). During hatching, membrane of egg pierced by egg tooth; crunching sound as bill is used as battering ram against shell; bird rotates in egg, cutting both membrane and shell until cap of shell is free.

Hatching asynchronous, but development period compressed such that chicks hatch in shorter interval than eggs laid; eggs laid over 5- to 6-d period generally hatch in 3–4 d (Dickerman and Gavino T. 1969). Generally hatch in order laid, but may hatch same day.

Parental assistance and disposal of eggshells. Adults usually toss eggshells over nest edge, but sometimes eggshells crushed in bottom of nest or pushed over edge by active chicks (Meyerriecks 1962).

YOUNG BIRDS

Condition at hatching. Semialtricial, with limited movement but eyes open.

MASS. Average weight of 62 hatchlings before first meal 16.0 g.

LINEAR MEASUREMENTS. At age 1 d, body length 95.3 mm, wing 15.9, bill 12.7, tarsus 15.9 (McVaugh 1975).

COLOR AND PLUMAGE PATTERN. At hatching, natal down wet and clinging but dry and fluffy in several hours. Down grayish brown forming crest on head, whiter on belly, less developed on wings (Gavino T. and Dickerman 1972). Down up to 19.1 mm long, darker gray on head.

COLOR OF BARE PARTS. Skin pink on forehead, crown, auricular area, and mouth lining, green around eyes; bill tipped black; tarsus and toes shades of greenish flesh (McVaugh 1975; see McVaugh's color illustrations and extensive descriptions of plumage and soft parts).

DEGREE OF COORDINATED MOVEMENT. Can only raise head for short periods at hatching, but able to move about nest later on day of hatching (Meyerriecks 1962).

RESPONSES, BODY ATTITUDE. Can only raise head for short periods, and eyes tend to close when head down (Douglass et al. 1965). In first day can

make *tik-tik-tik-tik* food-begging calls (Meyerriecks 1962).

Growth and development. Weight of 1 chick: 11.5 g (hatching), 16.5 (day 2), 88.8 (day 7), 132.2 (day 14), 173.2 (day 21). Reach minimal adult mass of 160 g in about 2 wk (Gavino T. and Dickerman 1969).

Chick weights after first day highly variable, partly because of differences in amount of food in crop or stomach. Gavino T. and Dickerman (1972) reported that sibling competition was not major factor in weight gain; even small chicks experienced normal growth, and differences leveled off after first week; small chick could be first, second, or last to hatch.

GROWTH OF BODY PARTS. Tarsus reaches 4.7 cm in 16–17 d; for daily increases in bill, tarsus, and eighth primary (see Gavino T. and Dickerman 1972).

MOLT INTO JUVENAL PLUMAGE. McVaugh (1975) reported that by day 3 upperparts mostly covered by gray down up to 19.1 mm long on forehead; by day 6, quills, with down tips, emerging from all tracts but crown and tail, and longest primary, secondary, and scapular pinfeathers 9.5 mm long; by day 8, crown and tail quills emerging; by day 11, most Juvenal feathers emerging; by day 13, quills and feathers covering most of chick; by day 31, body "well covered" with Juvenal feathers, but gray down still attached to some.

BEHAVIOR. At 9–11 d, parents begin to leave chicks alone during day, and chicks adopt Bittern Stance (see above, Behavior: social and interspecific behavior) when approached. By 3 wk, chicks will regurgitate food if disturbed. In nest will stretch wing over leg, preen, and head-scratch.

Meyerriecks (1962) reported first wing-flapping at 7–8 d, wing-preening by 6 d, fear response by 4–5 d (crouch, then Bittern Stance, then try to leave nest), climbing well by end of 1 wk, wing-stretching and forward stretching of head and neck by 8 d, snapping at flying insects by 9 d, jumping to nest from branch by 15 d, making short flights by 21 d.

LOCOMOTION. On day 3, chicks peep but cannot leave nest; on day 5, extend wings and give *haa* call; soon climb using wings, neck, beak, and feet; by day 15 can escape capture by climbing (Douglass et al. 1965). Cooley (1942) reported that by day 8–11, chicks left nest when approached, but quickly returned to nest when investigator entered blind. By 3 wk, chicks climb and move easily and will cling to branch with tenacity if handled. Will swim if they fall into water, with only top portion of back showing (Hindwood 1933).

PARENTAL CARE

Brooding. Both adults brood chicks. Meyerriecks (1962) reported that adults brood almost constantly during week 1, after that spend progressively less time until brooding ceases by end of week 3. On hot days adult may shade young, standing with wings spread, especially at exposed nests (Dickerman and Gavino T. 1969). Adults utter *cuck cuck* calls as they approach to brood or feed chicks (Cooley 1942).

Feeding. Food delivered to chicks by both parents. Chicks give food-begging call, low-pitched *tik-tik-tik-tik*, from first day. First fed at age 1 d. Usually fed soon after nest relief. First few days, well-digested food regurgitated directly into chick's mouth or onto nest platform. At age 1 wk, chicks grasp parent's bill crosswise and take food directly. By 2–3 wk, both parents may feed chicks at same time. Number of feedings/d increases during first and second week, then reduced to 2–3/d. Chicks fed at any time of day in New York colonies (Meyerriecks 1962). At age 6 d, chicks fed every 20 min, less frequently thereafter; after feeding, both adult and young preen, stretch, and rest (Cowles 1930 for *B. striatus*).

Food at first is nearly fluid; less-digested and larger items given to older chicks. Chicks fed in nest tree after they leave nest, and at other locations after they leave nest tree. Cooley (1942) reported that adult may feed all, some, or 1 of chicks at a particular feeding.

Nest sanitation. No attempt at nest sanitation by adults, but when strong enough to walk, chicks void over edge of nest (Cooley 1942). When a few days old, chicks shuffle backward to nest rim to void, later walk to rim; nest and surrounding area usually covered with excrement (Meyerriecks 1962).

Parental carrying of young. Not reported.

COOPERATIVE BREEDING

Not reported, and unlikely.

BROOD PARASITISM

Not reported, but 9- and 10-egg clutches reported (Huey 1926, Howsley 1936), suggesting egg dumping.

FLEDGLING STAGE

Departure from nest. At 16–17 d. Flying at 21–22 d. In a Michigan nest, young stayed in nest 15–16 d (Cooley 1942). Meyerriecks (1962) reported nestling period of 16–17 d and flight at 21–23 d. Gavino T. and Dickerman (1972) reported young flying up to 10 m at age 21–25 d. Young fly around nest tree while parents forage and fly to

nest tree when parents return to feed them 2–3 times/d. Young fly to feeding grounds with parents by 25 d.

IMMATURE STAGE

Time to independence unknown, but estimated at 30–35 d. Fledged young will enter water after prey and swim well. Most behavior at this stage little known. Hindwood (1933) reported for *B. striatus* that young, after leaving nest, stay in mangroves ≥ 1 mo where they are fed by adults and “taught” how to forage for fish. Observations of marked birds needed.

DEMOGRAPHY AND POPULATIONS

MEASURES OF BREEDING ACTIVITY

Age at first breeding; intervals between breeding. Usual age of first breeding probably 2 yr. At a New York colony, several breeding birds were in “immature” plumage, suggesting breeding at age 1 yr for some birds (Meyerriecks 1962). Undoubtedly attempts to breed every year, but data lacking.

Clutch. NUMBER OF EGGS. Usually 3–5, but variable among nests and colonies. Latitudinal differences not apparent. No statistical difference from first to second broods. Data: 4 or 5 eggs in N. Carolina (McVaugh 1975); in New York for 76 first clutches, 3 eggs ($n = 25$), 4 (38), 5 (11), 6 (2) (Meyerriecks 1962); in San Blas, Mexico, for 185 nests: 2 eggs ($n = 53$), 3 (125), 4 (7) (Dickerman and Gavino T. 1969). In Washington State, usually 4 or 5 eggs reported (Jewett et al. 1953). In Kentucky, mean of 4.3 eggs and young (Mengel 1965). Large clutches of 7 (Coffey 1981), 8 and 9 (Huey 1926), and 10 (Howsley 1936) reported, but not definitively laid by 1 individual.

NUMBER OF CLUTCHES/SEASON. Most individuals in northern latitudes (Meyerriecks 1962) probably lay 1 clutch; farther south (Dickerman and Gavino 1969), generally double brooded. In all areas may renest after failure of initial nesting (Meyerriecks 1960, 1962, Dickerman and Gavino T. 1969). In Mexico, 32 pairs laid normal second clutches, 1 pair 9 d after first brood fledged, and 7 nests had new eggs laid in same nest after egg destruction or abandonment (usually after 5–8 d), likely but not certainly laid by same pair (Dickerman and Gavino T. 1969).

Annual and lifetime reproductive success. Few data, none with marked birds. Widely dispersed and cryptic nature of solitary nesting pairs, together with small colony size of colonial nesters, makes collecting meaningful data on reproductive success difficult. Estimates that do not

correct for unobserved losses may produce spuriously high results (Erwin and Custer 1982).

In Missouri, 78.8% of 62 nests successful (Kaiser and Reid 1987). In 104 nests in San Blas, Mexico, 191 chicks fledged from 211 of 290 eggs (Dickerman and Gavino T. 1969). In 76 nests in New York, 60 had at least 1 egg hatch; in 51 nests, all eggs hatched, with 1 egg failure in 8 nests, and 2 in 1 nest (Meyerriecks 1962).

LIFE SPAN AND SURVIVORSHIP

Few data. Banded bird recovered in New Jersey age 4 yr (Cooke 1938) and in Mexico age 7 yr 11 mo (Clapp et al. 1982).

DISEASE AND BODY PARASITES

Disease. Green Herons are amazingly tolerant of mosquitoes, their only defensive movement being mild head-shaking (Edman et al. 1984); hence may be susceptible to mosquito-vectored diseases.

Body parasites. *Cercaria megalura* (larva of a species of *Philophthalmus*) has been found in orbits of Green Herons in s. Indiana (Fisher and West 1958). Other parasites: 2 cestodes, *Hymenolepis ardea* and *Ophiovalipora minuta*, in Indiana (Coil 1950); nematodes, *Strongyloides ardeae* in Louisiana and *S. cubaensis* in Cuba (Little 1966); nematodes *Tetrameres ardamericanus*, *Contra-caecum microcephalum*, and *Agamascaris* sp. in Massachusetts; also *T. ferminis* from Cuba (Boyd 1966); a blood parasite, *Leucocytozoon iowense*, in Iowa (Coatney 1938); a nasal mite, *Tinaminysus belopoliskii*, in Louisiana (Pence 1972).

CAUSES OF MORTALITY

Little known. Predation may be severe on eggs and chicks. Of 290 eggs at San Blas (Nayarit), Mexico, 24 infertile, 40 broken or disappeared (Dickerman and Gavino T. 1969). Kaiser and Reid (1987) found nest losses were primarily from predation, wind, and weather, also heat stress and early embryo death.

Few reports of mortality in adults. Few instances of predation or attempted predation reported, although species' choice of dense habitats reduces risk but also opportunities for observation. No evidence of adverse affects of competition from other species.

RANGE

Initial dispersal from natal site. Disperses from nesting site prior to fall migration. See Migration; also Distribution: winter range.

Fidelity to breeding site. Probable, but no data from marked birds. Some natal philopatry assumed, but birds banded as chicks in 1 area later seen far away. Evidence needed from marked birds.

Dispersal from breeding site or colony. Pronounced postbreeding dispersal, often into nonbreeding areas (see Migration) may be found wherever there is suitable foraging habitat within range.

POPULATION STATUS

Numbers. Because most Green Herons are solitary nesters and are dispersed widely through marine and freshwater habitats, estimating numbers is difficult. Also, colonies tend to be small. Species is not included in many surveys of colonially nesting waders (e.g., Spendelow and Patton 1988) because of sparse data, which if reported, might be misleading (J. A. Spendelow pers. comm.). Hence, few data on populations or colony sites exist.

Reported as present in 33 colonies in 7 East Coast states, with estimate (an underestimate) of 526 breeding adults (Custer and Osborn 1977). Historically, colonies of 100 pairs reported for Tampa Bay, FL, in 1912, and 400 from Orange Lake, FL, in 1916 (Howell 1932), but more recent colony data report lower numbers, e.g., 38 pairs from Alligator Bay, NC, in 1969 (Custer and Osborn 1977). In San Blas, Mexico, about 300 pairs reported for a 1964 colony in mangroves (Dickerman and Gavino T. 1969).

Trends. With poor data on numbers (see above), trends difficult to determine accurately. No indication of overall population decreases; suggestion of increases from available data. From 1968–1979, North American Breeding Bird Survey data indicated slight but significant increases in central and eastern regions, also for continent. No state showed a decline. Florida and Louisiana had highest means; increases noted for Alabama, California, Mississippi, Wisconsin, and Texas. Increases also noted for western portion of Great Lakes Plain, central and southern part of Upper Coastal Plain, and southern portion of Lower Coastal Plain (Robbins et al. 1986). Colony-based census data needed, however, as reported increases in some areas may reflect increased coverage (Broadbooks 1946).

Positive trend of long-term population expansion at northern and central range limits noted elsewhere.

POPULATION REGULATION

Poorly studied; needs attention. Habitat alteration (especially drainage of wetlands) affects breeding and foraging habitat, and hence potentially numbers. Storms can cause substantial mortality during nesting season (Wheelock 1906); likewise predation, especially at accessible nests. Island nesting probably helps deter predators

and thus boost reproduction, but data lacking. No data to suggest that interspecific competition is a major regulator of populations.

CONSERVATION AND MANAGEMENT

EFFECTS OF HUMAN ACTIVITY

Shooting and trapping. In past often shot for food (Bent 1926); P. A. Taverner (in McAtee 1919) reported that Green Heron was "very good, a little more delicate than the Night Heron." Historically, Green Heron subject to predator-control programs at fish hatcheries (Slipp 1944); probably continues today at some hatcheries.

Pesticides and other contaminants. Found to accumulate persistent pesticides, but no evidence of general reproductive failure, despite some localized effects.

Except for DDE, mirex, and *cis*-nonachlor, frequency of organochlorines in Green Heron eggs "considerably" lower than average for 17 species of herons, bitterns, ibises, and storks from eastern U.S. (Ohlendorf et al. 1977, 1979). Mirex particularly common in eggs from Savannah National Wildlife Refuge (NWR), SC, where highest concentration of DDE was also found (1.4 parts per million [ppm] wet weight). PCBs usually less than half the average for all species, but highest in birds from Potomac River, MD (1.6 ppm). From Southern Atlantic, inland, and Gulf Coast regions combined, Green Heron egg ($n = 89$) percentages with detectable residues were: DDE (97%), DDD (1%), DDT (7%), dieldrin (10%), mirex (8%), Oxychlorane (1%), *cis*-chlorane (1%), *cis*-nonachlor (1%), PCBs (28%).

Eggshell thinning evident in comparisons with pre-1947 egg samples, but not considered deleterious (see Breeding: eggs). In Tennessee Valley, DDE concentrations fairly high (detected in 100% of eggs sampled; mean = 3.9 ppm wet weight; range = 1.3–12.0 ppm; $n = 8$).

White et al. (1988) reported that DDE contamination of Green Heron eggs at Yazoo NWR, MS, in 1984–1985 ranged from 0.63 to 43.0 ppm wet weight ($n = 59$, mean = 7.4). DDE concentrations correlated negatively and significantly with hatching success; threshold concentrations = 5.1–10.0 ppm. Sources of contamination, although probably local, remain unknown.

Disturbance. Increased recreational use of river channels leads to decreased use by Green Herons and reduced foraging time, but does not affect heron use of backwater habitat such as oxbow ponds, side channels, and marginal pools (Kaiser and Fritzell 1984).

Human/research impacts. In Mexico, investigator disturbance, particularly during early egg-laying period, caused nest abandonment; also mortality through egg breakage and exposure to sun (Dickerman and Gavino T. 1969).

MANAGEMENT

Conservation status. No population considered to be threatened.

Measures proposed and taken. Because threat of pesticides to reproductive efforts of Green Herons appears to be minor, and because most populations are stable or increasing, management has not been major focus.

Primary concern is conservation and management of wetlands of all sizes, including small ones, as these are habitats on which Green Herons depend. Various parks, refuges, lakes, and recreation areas managed by various government agencies support Green Heron populations across North America. Management of this species should include (1) identifying wetlands and especially protected areas used by Green Herons and (2) including species' needs in habitat management plans for areas.

Some developments have enhanced habitat for this species: e.g., reservoirs that create permanent water in otherwise dry environments, permanent water marshes created along coast for mosquito control, and wetland management for waterfowl. Dredged material islands, created for channels, are widely used by colonially nesting wading birds; Green Herons have been reported nesting on dredge material islands in Texas, Florida, N. Carolina, New Jersey, and Great Lakes region (Landin and Soots 1977).

APPEARANCE

MOLTS AND PLUMAGES

Molts poorly known, based on Bent 1926, Meyerriecks 1962, McVaugh

Despite its widespread geographic distribution and abundance the molts and plumages of this species are poorly known. In addition, Cramp's (1977) account of the sequence of molts in the closely related Striated Heron is confused and incorrect. No specific studies on Prebasic and Prealternate molts have been done in either species. Wheelock (1906), Hindwood (1933), Gavino T. and Dickerman (1972), and McVaugh (1975) documented the development of plumage and bare part colors of young birds in detail from hatching to fledging.

Hatchlings. Downy. At hatching, down on head, neck, and wings. Down longest (up to

Table 1. Average mean measurements (mm) of Green Herons *B. v. virescens* and *B. v. anthonyi* (after Oberholser 1912). Sample size (*n*) shown as male/female.

| | <i>B. v. virescens</i> <i>n</i> = 14 / 13 | <i>B. v. anthonyi</i> <i>n</i> = 11 / 9 |
|----------------|--|--|
| WING | | |
| Male | 181.1 | 196.3 |
| Female | 175.1 | 189.9 |
| TAIL | | |
| Male | 66.5 | 72.9 |
| Female | 65.5 | 70.3 |
| EXPOSED CULMEN | | |
| Male | 60.9 | 60.8 |
| Female | 59.8 | 60.3 |
| TARSUS | | |
| Male | 53.0 | 53.3 |
| Female | 51.2 | 52.6 |

19.1 mm) on forehead and crown, forming a crest (McVaugh 1975). Additional down appears on back, neck, and femoral tract by 3 d; down darkest on head, its color described as grayish brown (Gavino T. and Dickerman 1972), hair brown (Bent 1926) to smoke gray (Palmer 1962); down shorter and densest on back, paler (light gray to nearly white) on underside (Bent 1926, Gavino T. and Dickerman 1972, McVaugh 1975).

Juvenal plumage. Prejuvenal molt complete. Juvenal pin-feathers, including primaries and secondaries, first appear in the alar, humeral, and interscapular region of dorsal tract by 4 d; feathers emerge on spinal, ventral, and femoral tracts by 5–6 d; on the neck by 7 d; on head (forehead, crown, and nape) and tail (rectrices) by 10–11 d. By 25 d, entire body covered with Juvenal plumage, down remaining only on head. All traces of down lost by 37 d (Gavino T. and Dickerman 1972, McVaugh 1975).

Contrary to some other authors (e.g., Palmer 1962), Bent (1926) and Oberholser (1974) state that sexes are distinguishable in Juvenal plumage. Ostensibly, in males, forehead and crown greenish black; sides of head and neck chestnut; malar stripe buffy white bordered above by blackish streak and below by blackish brown streaking which extends down side of neck; sides of neck yellowish white broadly streaked with diffuse dark chestnut streaks with ventral portion of throat and neck largely whitish with some

narrow dark chestnut streaking medially; scapulars and back dark greenish gray with brownish cast; wing coverts are dark greenish gray except lesser coverts are edged chestnut, and median and greater coverts are rounded at tips, not pointed as in adults, with pale buff edgings and a triangular buffy white spot at the tip of each rachis; these spots rapidly fade to white and wear away. Lesser coverts dark gray edged chestnut; wing lining variegated. Remiges and rectrices dark gray with greenish cast; secondaries and primaries tipped with white in decreasing amounts from the inner secondaries to the outer primaries; underparts buffy white, streaked with buff, brown, or dusky; undertail coverts white tinged with buff.

Bent and Oberholser described females as differing from males in having chestnut streaks on crown, and having sides of head and neck streaked with chestnut, buff and dusky. Very few examples of sexual dimorphism in Juvenal plumage exist. Therefore, further studies should evaluate the validity of this purported early sexual dimorphism.

Basic I plumage. Most birds replace all body plumage (i.e., not including flight feathers) during the Prebasic I molt, although some late-hatching birds undergo little or no molt (Townsend *in* Bent 1926). Occurs from few weeks (at most) after attaining flight until late winter (Fig. 5). Oberholser (1974) erroneously states that Basic I plumage is attained by wear rather than by molt.

Plumage similar to Juvenal plumage but forehead and crown dark greenish gray, clear white on chin, throat, and underparts; upper breast and neck more broadly striped.

Alternate I plumage. Prealternate I molt partial, limited to "taking on of some additional plumes on the back" as well as "a few new feathers...on the shoulders." Most birds in this plumage have not acquired plumes as long as adults of the same sex in Definitive Alternate plumage (Townsend *in* Bent 1926). Molt occurs late winter or early spring (Palmer 1962). Extent probably more limited than in subsequent Prealternate molts (Palmer 1962).

Plumage similar to Definitive Alternate but upperparts duller; inner primaries, outer secondaries, primary coverts, and greater coverts with white or buffy tips; primaries duller; greater coverts more rounded; more white on lower throat and breast; and plumes in both sexes shorter (male plumes longer than female) (Palmer 1962, Oberholser 1974).

Basic II plumage. Prebasic II molt usually complete, but some birds may retain at least some Juvenal secondaries (e.g., University

Washington specimens [45754, 35962, 45348]; Palmer 1962). Molt begins early summer, earlier than subsequent Prebasic molts, and is completed by Nov (Palmer 1962: Fig. 5).

Plumage generally similar to Definitive Basic. Crown glossy dark greenish gray or bottle green, with elongated erectile feathers; sides of head and most of neck chestnut or maroon; white stripe down throat; spots on lower throat and breast elongated and narrower than in earlier plumages; upperparts mostly glossy dark greenish gray; underparts brownish gray; wings dusky gray-green; buff edges to all coverts.

Definitive Alternate plumage. Definitive Prealternate II molt more extensive than first Prealternate molt involving replacement of head and neck feathers, including ornamental plumes, and some body plumage (Palmer 1962).

Forehead, crown, and lower border of eye dark grayish bottle green with medium length erectile plumes extending from rear of crown forming a bushy crest when raised; remainder of head, neck, breast, and upper sides purplish maroon becoming mouse gray on remainder of underparts; thighs rufescent.

Malar stripe cream buff bordered above by blackish streak; chin and throat buffy white, medial portion of lower neck and breast white, spotted on upper neck and streaked on breast with grayish with variegated white and brownish streaks extending to the upper belly. Upper back and scapulars covered with medium length bluish gray plumes with a greenish cast; lower back greenish blue-gray; tail dark greenish gray with bluish cast. Outer scapulars, lesser and median coverts dark greenish gray margined with tawny buff, remainder of wing dark bluish gray with greenish cast, outer webs of tertials and innermost secondaries edged with white; leading edge of bend of wing buffy white; lining of wing slate gray margined with buff (Oberholser 1974). Male more brightly colored than female.

Definitive Basic plumage. Definitive Prebasic Molt complete, from late summer or fall through winter. Plumage similar to Definitive Basic except plumes shorter.

Polymorphism. Unusual plumages include erythristic individuals from Cuba and Panama (Meyerrieks 1962). Head and neck reddish chestnut with black cap, no white on throat and foreneck, wing coverts narrowly edged rusty, young nearly uniform brown.

BARE PARTS

McVaugh (1975) provides the most detailed account of bare part colors in nestlings, and is summarized below unless stated otherwise. Other

less detailed accounts that differ in some respects from McVaugh are Wheelock (1906), Hindwood (1933), and Gavino T. and Dickerman (1972).

Bill. At hatching, upper mandible pale greenish gray at base, culmen dark pinkish gray proximally and black distally, tip black; rounded sides of bill pinkish; black line runs just above commissural point to about one half length of bill; lower mandible black distally with ivory tip; by 3 d, upper mandible yellowish pink, black at tip, distal one half of tomlia black; sides of lower mandible yellowish at base becoming pinkish and then blackish distally with small ivory tip. By 11 d, upper mandible greenish orange, more greenish basally, culmen pale orange with black tip, tomlia black distally; lower mandible greenish orange basally becoming black distally with small ivory tip. By 24 d, upper mandible orange, distal end black with ivory tip; lower mandible orange, paler distally with small black tip. In adult, bill brownish black, lower mandible dusky green, yellowish at base; bill glossy black in breeding season. Mouth lining pale pinkish from hatching through fledging.

Iris. At hatching pale gray (off white in some birds) through 8 d becoming grayish white from 11 through 13 d, greenish to yellowish white from 17 d through fledging. In adults, orange and/or yellow, but may be deep orange during intense display.

Bare skin on head. At day 6, skin of forehead and crown pinkish, auricular area green, body pale yellow-green to gray-green; lores yellow to greenish yellow from 5 d through fledging (Gavino T. and Dickerman 1972, McVaugh 1975). Eye-ring pale yellow-green at 3 d, pale green from 6 through 13 d, greenish yellow by 17 d, becoming greenish blue in some birds between 24 d and fledging. In adult, upper portion of lores and bare skin around upper eye and posterior lower portion of eye dull yellow-green, lower portion of lores blackish; lores bluish black at height of breeding season.

Legs and feet. At hatching, tibiotarsus greenish flesh to flesh, heel and tarsometatarsus pale flesh; toes flesh above, pale flesh below, nails ivory. By 3 d, tarsi and toes pale yellow; nails pale bluish gray with paler tips. By 11 d, tibiotarsus pale green to yellow-green, heel and tarsometatarsus yellow; toes darker green than tibiotarsus with deep yellow undersides; nails medium gray with pale tips. From 17 d through fledging, tarsometatarsus yellow, pale yellow, or pale orange-yellow, toes darker greenish yellow above and orange beneath; web between second and third toes orange to orange-green; nails medium horn

with paler tips. Tibiotarsus pale yellowish green (some grayish green) from 24 d through fledging. Adult, yellow or orange-yellow, glossy orange in breeding season.

MEASUREMENTS

LINEAR

Bill lengths for 16 adults averaged 6.3 cm and 5.9 cm (Niethammer and Kaiser 1983). Females average larger in some parts of range, smaller in others (Oberholser 1912). See Table 1 for measurements of males and females of *B. v. virescens* and *B. v. anthonyi* (and see Oberholser 1912 for measurements throughout geographic range).

MASS

Weights for 16 adults averaged 241 g, for 14 immatures 219 g (Niethammer and Kaiser 1983).

PRIORITIES FOR FUTURE RESEARCH

1. Get better information on wintering areas of specific North America populations and their migration pathways. This may require additional attention or banding studies.

2. Develop better understanding of habitat use and choice, especially on migration and in wintering areas.

3. Factors influencing reproductive success and population regulation in this species remain poorly known and demand attention.

4. Molts and plumages are clearly in need of study. Correct aging of individuals depends on such knowledge.

5. Systematics of species remains unresolved. A better understanding of population structure of *B. virescens*, and also of Neotropical *B. striatus*, is needed. Future research should focus on anatomical and biochemical studies of the several populations. This research is of interest from both a systematic and conservation perspective.

6. Flexibility of species in nesting colonially or solitarily in the same areas deserves additional attention.

7. Biology of populations on islands, where species can be quite common, remains little known. If Galapagos Heron is any indication, such studies should prove interesting and affect thinking about breadth of adaptability of species.

8. Suggestions that recreational disturbance adversely affects habitat use should be pursued in additional areas. Extent that disturbance adversely affects this rather retiring species is

critical for proper management of populations on refuges and reserves where human recreation is encouraged.

ACKNOWLEDGMENTS

Danny Bystrak of the Bird Banding Laboratory provided banding recovery data. Sandra L. L. Gaunt of the Borror Laboratory of Bioacoustics at Ohio State University provided the spectrograph. Lloyd Kiff of the Western Foundation of Vertebrate Zoology provided nest and egg data. T. Custer reviewed portions of this manuscript and made helpful suggestions. Cover photo © Arthur and Elaine Morris/VIREO.

REFERENCES

- Adams, L. W., L. E. Dove, and T. M. Franklin. 1985. Use of urban stormwater control impoundments by wetland birds. *Wilson Bull.* 97: 120–122.
- Amadon, D. 1953. Avian systematics and evolution in the Gulf of Guinea, the J. G. Correia collection. *Bull. Am. Mus. Nat. Hist.* 100: 393–452.
- American Ornithologists' Union. 1976. Thirty-third supplement to the AOU Check-list. *Auk* 93: 875–879.
- American Ornithologists' Union. 1983. Check-list of North American birds, 6th ed. Am. Ornithol. Union, Washington, D.C.
- American Ornithologists' Union. 1993. Thirty-ninth supplement to the American Ornithologists' Union Check-list of North American Birds. *Auk* 110: 675–682.
- Avakian, A. 1984. Green-backed Heron spends two consecutive winters on pond in northwestern South Carolina. *Chat* 48: 13–14.
- Barker, S. H. 1901. Does the Green Heron fish in deep water? *Bird Lore* 3: 141.
- Baynard, O. E. 1912. Food of herons and ibises. *Wilson Bull.* 24: 167–169.
- Becker, C. N., and S. M. Byers. 1976. Green Heron as a potential prey species of the Broad-winged Hawk. *Bird Banding* 47: 365–366.
- Bent, A. C. 1926. Life histories of North American marsh birds. U.S. Natl. Mus. Bull. No. 135.
- Betts, B. J., and D. L. Betts. 1977. The relation of hunting site changes to hunting success in Green Herons and Green Kingfishers. *Condor* 79: 269–271.
- Blake, C. H. 1948. More data on the wing flapping rates of birds. *Condor* 50: 148–151.
- Bock, W. J. 1956. A generic review of the family Ardeidae (Aves). *Am. Mus. Novit.* No. 1779.
- Boyd, E. M. 1966. Observations on nematodes in North America including three new species and new host and state records. *J. Parasitol.* 52: 503–511.
- Broadbooks, H. E. 1946. Anthony Green Heron at Yaquina Bay, Oregon. *Murrelet* 27: 12.
- Brooks, W. S. 1923. An interesting adaptation. *Auk* 40: 121–122.
- Brown, L. H., E. K. Urban, and K. Newman. 1982. The birds of Africa, vol. I. Academic Press, New York.
- Bryant, H. C. 1914. Birds as destroyers of grasshoppers in California. *Auk* 31: 168–177.
- Bull, J. 1974. Birds of New York State. Cornell Univ. Press, Ithaca, NY.
- Burger, J. 1978. The pattern and mechanism of nesting in mixed-species heronries. Pp. 45–58 in *Wading birds* (A. Sprunt IV, J. C. Ogden, and S. Winkler, eds.). Nat. Aud. Soc. Res. Rep. No. 7, New York.
- Burger, J., J. Shisler, and F. H. Lesser. 1982. Avian utilization on six salt marshes in New Jersey. *Biol. Conserv.* 23: 187–212.
- Burns, F. L. 1917. Comparative periods of deposition and incubation of some North American birds. *Wilson Bull.* 27: 275–286.
- Caccamise, D. F., and J. Fischl. 1985. Patterns of association of secondary species in roosts of European Starlings and Common Grackles. *Wilson Bull.* 97: 173–182.
- Caldwell, G. S., and R. W. Rubinoff. 1983. Avoidance of venomous sea snakes by naive herons and egrets. *Auk* 100: 195–198.
- Campbell, K. E., Jr. 1976. An early avifauna from Haile XVA, Florida. *Wilson Bull.* 88: 345–347.
- Clapp, R. B., M. K. Klimkiewicz, and J. H. Kennard. 1982. Longevity records of North American birds: Gaviidae through Alcidae. *J. Field Ornithol.* 53: 81–208.
- Clarke, J. A., B. A. Harrington, T. Hruby, and F. E. Wasserman. 1984. The effect of ditching for mosquito control on salt marsh use by birds in Rowley, Massachusetts. *J. Field Ornithol.* 55: 160–180.
- Coatney, G. R. 1938. Some blood parasites from birds of the Lake Okaboji region. *Am. Mid. Nat.* 20: 336–340.
- Coffey, B. B., Jr. 1981. A past Green heronry colony in Memphis. *Migrant* 52: 44.
- Coil, W. H. 1950. The genus *Ophiovalipora* HSU (Cestoda: Dilepididae) with the description of *Ophiovalipora minuta* Sp. Nov. from the Green Heron (*Butorides virescens* L.). *J. Parasitol.* 36: 55–61.
- Cooke, M. T. 1938. Returns of banded birds: recoveries of banded marsh birds. *Bird Banding* 9: 80–87.
- Cooley, E. G. 1942. The nesting of a pair of Green Herons. *Jack-Pine Warbler* 20: 3–9.
- Cowles, R. B. 1930. Notes on the nesting of the African Green Heron (*Butorides atricapilla*) in Natal. *Auk* 47: 465–470.
- Cramp, S., ed. 1977. Handbook of the birds of Europe, the Middle East, and North Africa: the birds of the Western Palearctic, vol. 1. Oxford Univ. Press, Oxford.
- Curry-Lindahl, K. 1971. Systematic relationships in herons (Ardeidae), based on comparative studies of behaviour and ecology. *Ostrich Sup.* 9.
- Custer, T. W., and R. G. Osborn. 1977. Wading birds as biological indicators: 1975 colony survey. U.S. Fish

- and Wildl. Serv. Spec. Sci. Rep. Wildl. No. 206, Washington, D.C.
- Davis, W. E., Jr. 1983. Observations of foraging behaviour of a Mangrove Heron. *Australian Birds* 18: 9.
- Debenedictis, P. A. 1978. Gleanings from the technical literature. *Birding* 10: 36-37.
- Debenedictis, P. A. 1994. Gleanings from the technical literature: 39th supplement to the AOU check-list. *Birding* 26: 59-61.
- del Hoyo, J., A. Elliott, and J. Stargatal, eds. 1992. *Handbook of the birds of the world*, vol. 1. Lynx Edicions, Barcelona.
- DeMay, I. S. 1942. An avifauna from Indian kitchen middens at Buena Vista Lake, California. *Condor* 44: 228-230.
- Dickerman, R. W., and G. Gavino T. 1969. Studies of a nesting colony of Green Herons at San Blas, Nayarit, Mexico. *Living Bird* 8: 95-111.
- Douglass, L. E., M. L. Bierly, and K. A. Goodpasture. 1965. Green Herons nest at Basin Spring. *Migrant* 36: 76-80.
- Eddy, G. 1952. Further notes on the Green Heron advance in Washington State. *Murrelet* 32 (1): 12.
- Edman, J. D., J. F. Day, and E. D. Walker. 1984. Field confirmation of laboratory observations on the differential antimosquito behavior of herons. *Condor* 86: 91-92.
- Erwin, R. M., and T. W. Custer. 1982. Estimating reproductive success in colonial waterbirds: an evaluation. *Colon. Waterbirds* 5: 49-56.
- Fisher, F. M., and A. F. West. 1958. *Cercaria megalura* Cort, 1914, the larva of a species of *Philophthalmus*. *J. Parasitol.* 44: 648.
- Fleming, W. J., B. P. Pullin, and D. M. Swineford. 1984. Population trends and environmental contaminants in herons in Tennessee Valley, 1980-81. *Colon. Waterbirds* 7: 63-73.
- Gaston, G. R., and P. G. Johnson. 1977. Nesting success and mortality of nestlings in a coastal Alabama heron-egret colony. *Northeast Gulf Science* 1: 14-22.
- Gavino T., G., and R. W. Dickerman. 1972. Nestling development of Green Herons at San Blas, Nayarit, Mexico. *Condor* 74: 72-79.
- Gordon, D. C. 1961. Herons of New York State. *Kingbird* 11: 11-18.
- Graf, W. 1946. Notes on the distribution and habits of Anthony Green Heron. *Murrelet* 27: 50-51.
- Griffie, W. E. 1954. Some Oregon nesting records. *Murrelet* 35: 48-49.
- Griscom, L. 1923. *Birds of the New York City region*. Handbook Series, No. 9, American Museum of Natural History, New York.
- Hancock, J., and J. Kushlan. 1984. *The herons handbook*. Harper and Row, New York.
- Hauser, D. C. 1957. Some observations on sun-bathing in birds. *Wilson Bull.* 69: 78-90.
- Hawbecker, A. C. 1949. Green Heron feeds on goldfish. *Auk* 66: 78-79.
- Higuchi, H. 1986. Bait-fishing by the Green-backed Heron *Ardeola striata* in Japan. *Ibis* 128: 285-290.
- Higuchi, H. 1988. Individual differences in bait-tasting by the Green-backed Heron *Ardeola striata* associated with territory quality. *Ibis* 130: 39-44.
- Hindwood, K. A. 1933. The Green-backed Mangrove Heron, pts. 1 and 2. *Emu* 33: 27-43; 97-102.
- Howard, H. 1936. Further studies upon the birds of the Pleistocene of Rancho La Brea. *Condor* 38: 32-36.
- Howell, A. H. 1932. *Florida bird life*. Fla. Dept. Fish and Game, Tallahassee.
- Howsley, L. B. 1936. Unusual sets of Bush-tit and Green Heron. *Condor* 38: 39.
- Hoyt, S. F. 1961. Foot-stirring in the Green Heron. *Wilson Bull.* 73: 386.
- Huey, L. M. 1915. Random notes from San Diego. *Condor* 17: 59-60.
- Huey, L. M. 1926. A prolific Anthony Green Heron. *Condor* 28: 94-96.
- Jewett, S. G. 1939. Anthony Green Heron in the state of Washington. *Condor* 41: 33.
- Jewett, S. G., W. P. Taylor, W. T. Shaw, and J. W. Aldrich. 1953. *Birds of Washington State*. Univ. Washington Press, Seattle.
- Junor, F. J. R. 1972. Estimation of the daily food intake of piscivorous birds. *Ostrich* 43: 193-205.
- Kaiser, M. S., and E. K. Fritzell. 1984. Effects of river recreationists on Green-backed Heron behavior. *J. Wildl. Manage.* 48: 561-567.
- Kaiser, M. S., and F. A. Reid. 1987. A comparison of Green-backed Heron nesting in two freshwater ecosystems. *Colon. Waterbirds* 10: 78-83.
- Kalter, L. B. 1932. A comparatively tame Eastern Green Heron. *Auk* 49: 342.
- Keenan, W. J. III. 1981. Green Heron fishing with mayflies. *Chat* 45: 41.
- Kramer, D. L., D. Manley, and R. Bourgeois. 1983. The effect of respiratory mode and oxygen concentration on the risk of aerial predation in fishes. *Can. J. Zool.* 61: 653-665.
- Kushlan, J. A. 1976. Feeding behavior of North American herons. *Auk* 93: 86-94.
- Kushlan, J. A. 1978. Feeding ecology of wading birds. Pp. 249-297 in *Wading birds* (A. Sprunt, IV, J. C. Ogden, and S. Winkler, eds.). Nat. Aud. Soc. Res. Rep. No. 7, New York.
- Kushlan, J. A. 1981. Resource use strategies of wading birds. *Wilson Bull.* 93: 145-163.
- Landin, M. C., and R. F. Soots. 1977. Colonial bird use of dredged material islands: a national perspective. *Proc. Colon. Waterbird Group* 1: 62-72.
- Larrison, E. J. 1940. The Anthony Green Heron in the state of Washington. *Murrelet* 21 (1): 1-3.
- Larrison, E. J. 1947. Present status of the Green Heron in Washington. *Condor* 49: 87.
- Laughlin, S. B., and D. P. Kibbe. 1985. *The atlas of breeding birds of Vermont*. Vermont Institute of Natural Sciences and Univ. Press of New England, Hanover, NH.
- Little, M. D. 1966. Seven new species of *Strongyloides* (Nematoda) from Louisiana. *J. Parasitol.* 52: 85-97.
- Lovell, H. B. 1958. Baiting of fish by a Green Heron. *Wilson Bull.* 70: 280-281.
- Marra, P., and M. Bull. 1986. The Norwalk Island heron

- colonies—a history. Connecticut Warbler 6: 23–26.
- McAtee, W. L. 1919. Further notes on the “fishy” flavor of birds. Auk 36: 100–101.
- McLean, D. D. 1930. The speed of flight in certain birds. Gull 12 (3): 1–2.
- McNeil, R., P. Drapeau, and R. Pierotti. 1993. Nocturnality in colonial waterbirds: occurrence, special adaptations, and suspected benefits. Curr. Ornithol. 10: 187–246.
- McVaugh, W., Jr. 1975. The development of four North American herons. II. Living Bird 14: 163–183.
- Mengel, R. M. 1965. The birds of Kentucky. Ornithol. Monogr. No. 3, Am. Ornithol. Union, Washington, D.C.
- Meyerriecks, A. J. 1960. Comparative breeding behavior of four species of North American herons. Publ. Nuttall Ornithol. Club No. 2, Cambridge, MA.
- Meyerriecks, A. J. 1962. Green Heron. Pp. 419–426 in Handbook of North American birds, vol. 1 (R. S. Palmer, ed.). Yale Univ. Press, New Haven, CT.
- Meyerriecks, A. J. 1966. Additional observations on “foot-stirring” feeding behavior in herons. Auk 83: 471–472.
- Meyerriecks, A. J. 1971. Further observations on use of the feet by foraging herons. Wilson Bull. 83: 435–438.
- Miller, A. H. 1945. Further records of birds from central California. Condor 47: 217–218.
- Monroe, B. L., Jr., and M. R. Browning. 1992. A re-analysis of *Butorides*. Bull. Br. Ornithol. Club 112: 81–85.
- Mousley, H. 1945. The nesting of the Eastern Green Heron, *Butorides virescens virescens* (Linn.) in the province of Quebec, May, 1942. Can. Field-Nat. 59: 51–52.
- Nickell, W. P. 1966. The nesting of the Black-crowned Night Heron and its associates. Jack-Pine Warbler 44: 130–139.
- Niethammer, K. R., and M. S. Kaiser. 1983. Late summer food habits of three heron species in northeastern Louisiana. Colon. Waterbirds 6: 148–153.
- Norris, D. 1975. Green Heron (*Butorides virescens*) uses feather lure for fishing. Am. Birds 29: 652–654.
- Oberholser, H. C. 1912. A revision of the subspecies of the Green Heron (*Butorides virescens* [Linnaeus]). Proc. U.S. Natl. Mus. 42: 529–577.
- Oberholser, H. C. 1974. The bird life of Texas, vol. 2. University of Texas Press, Austin.
- Ohlendorf, H. M., E. E. Klaas, and T. E. Kaiser. 1977. Organochlorine residues and eggshell thinning in anhingas and waders. Proc. Colon. Waterbird Group 1: 185–195.
- Ohlendorf, H. M., E. E. Klaas, and T. E. Kaiser. 1979. Environmental pollutants and eggshell thickness: anhingas and wading birds in the eastern United States. U.S. Fish and Wildl. Serv. Spec. Sci. Rep. Wildl. No. 216, Washington, D.C.
- Osborn, R. G., and T. W. Custer. 1978. Herons and their allies: atlas of Atlantic Coast colonies, 1975 and 1976. Biological Services Program, U.S. Fish and Wildl. Serv. FWS/OBS-77/08.
- Paton, P., and L. H. MacIvor. 1983. Green Heron in Hawaii. Am. Birds 37: 232–233.
- Payne, R. B. 1974. Species limits and variation of the New World Green Herons *Butorides virescens* and Striated Herons *B. striatus*. Bull. Br. Ornithol. Club. 94: 81–88.
- Peck, G. K. and R. D. James. 1983. Breeding birds of Ontario: nidiology and distribution. Royal Ontario Museum, Toronto.
- Pence, D. B. 1972. The nasal mites of birds from Louisiana. I. Dermanyssids (Rhinonyssinae) from shore and marsh birds. J. Parasitol. 58: 153–168.
- Peterjohn, B. G., and W. Zimmerman. 1989. The birds of Ohio. Indiana Univ. Press, Bloomington.
- Peters, J. L. 1979. Check-list of birds of the world, vol 1, 2nd ed. (E. Mayr and G. W. Cottrell, eds.). Museum of Comparative Zoology, Cambridge, MA.
- Peterson, R. T. 1980. A field guide to the birds, 4th ed. Houghton Mifflin, Boston, MA.
- Peterson, R. T. 1990. A field guide to western birds, 3d ed. Houghton Mifflin, Boston, MA.
- Recher, H. F., and J. A. Recher. 1980. Why are there different kinds of herons? Trans. Linn. Soc. NY 9: 135–158.
- Recher, H. F., R. T. Holmes, W. E. Davis, Jr., and S. Morton. 1983. Foraging behavior of Australian herons. Colon. Waterbirds 6: 1–10.
- Ridgway, R. 1912. Color standards and color nomenclature. Privately printed, Washington, D.C.
- Robbins, C. S., D. Bystrak, and P. H. Geissler. 1986. The breeding bird survey: its first fifteen years, 1965–1979. U.S. Fish and Wildl. Serv. Res. Publ. 157, Washington, D.C.
- Robbins, M. B., and D. A. Easterla. 1992. Birds of Missouri: distribution and abundance. Univ. Missouri Press, Columbia.
- Ryder, R. A. 1979. Status, distribution, and movements of ciconiiforms in Colorado. Proc. Colon. Waterbird Group 3: 49–57.
- Sandilands, A. P. 1977. Green Heron nesting in a Wood Duck box. Auk 94: 390.
- Scott, W. E. D. 1890. On birds observed at the Dry Tortugas during parts of March and April, 1890. Auk 7: 301–314.
- Seibert, H. C. 1951. Light intensity and the roosting flight of herons in New Jersey. Auk 68: 63–74.
- Sheldon, F. H. 1987. Phylogeny of herons estimated from DNA–DNA hybridization data. Auk 104: 97–108.
- Sibley, C. S., and B. L. Monroe, Jr. 1990. Distribution and taxonomy of birds of the world. Yale Univ. Press, New Haven, CT.
- Slipp, J. W. 1942. Green Heron at Tacoma, Washington. Condor 44: 35–36.
- Slipp, J. W. 1944. Additional records of the Green Heron in the Puget Sound basin. Condor 46: 35–36.
- Spendelow, J. A., and S. R. Patton. 1988. National atlas of coastal waterbird colonies in the contiguous United States: 1976–82. U.S. Fish and Wildl. Serv. Biol. Rep. No. 88 (5).
- Steirly, C. C. 1964. Green Heron nesting in loblolly pine plantation. Raven 35: 51.

- Townsend, C. W. 1928. The song of the Green Heron (*Butoroides* [sic] *virescens virescens*). Auk 45: 498-499.
- Wetmore, A. 1931. The avifauna of the Pleistocene in Florida. Smithson. Misc. Coll. 85 (2): 1-41.
- Wetmore, A. 1965. The birds of the republic of Panama. Part 1. Tinamidae (tinamous) to Rynchopidae (skimmers). Smithson. Misc. Collect. 150.
- Wheelock, I. G. 1906. Nesting habits of the Green Heron. Auk 23: 432-436.
- White, D. H., W. J. Fleming, and K. L. Ensor. 1988. Pesticide contamination and hatching success of waterbirds in Mississippi. J. Wildl. Manage. 52: 724-729.
- Wood, H. B. 1933. Flight speeds of some birds. Auk 50: 452-453.

ABOUT THE AUTHORS

William E. Davis, Jr., is professor and past chairman of the Division of Science and Mathematics, College of General Studies, Boston University. He has studied herons in North and South America and in Australia. He is past president of the Nuttall Ornithological Club and Association of Field Ornithologists. He is currently president of Bird Observer of Eastern Massachusetts, Inc., and serves on the councils of the Colonial Waterbird and Wilson Ornithological societies. He is interested in the history of ornithology and has published the books History of the Nuttall Ornithological Club, 1873-1986; Dean of the Birdwatchers: A Biography of Ludlow Griscom; and Contributions to the History of North American Ornithology (edited with Jerome Jackson). Current address: Division of Science and Mathematics, College of General Studies, Boston University, Boston MA 02215.

James A. Kushlan has studied colonial waterbirds and their wetland habitats extensively. He is the author of more than 150 papers, mostly on wetlands and wetland species. He also is coauthor of monographic books on the biology of several groups of colonial waterbirds (The Herons Handbook and Storks, Ibises, and Spoonbills of the World) and is coeditor of the forthcoming book Heron Conservation. He also is the president-elect of the Colonial Waterbird Society and former editor of the journal Colonial Waterbirds. He is associate editor of the journal Wetlands, a fellow and council member of the American Ornithologists' Union, a member of the Steering Committee of the International Wetland and Waterfowl Research Bureau, and cochair of the Herons Specialist Group. Current address: Department of Biology, The University of Mississippi, University, MS 38677.

Appendix. Stomach contents of 27 Green Herons from nw. Louisiana in late summer. A = percent frequency of stomachs with prey type; B = percent of total diet by weight. After Niethammer and Kaiser (1983).

| | A | B |
|--|----|-----|
| FISH | 93 | 93 |
| Mosquitofish (<i>Gambusia affinis</i>) | 11 | 1 |
| Shiners (<i>Notropis</i> spp.) | 7 | 2 |
| Sunfish (<i>Lepomis</i> spp.) | 26 | 35 |
| Pirate perch (<i>Aphredoderus sayanus</i>) | 4 | 2 |
| Threadfin shad (<i>Dorosoma petenense</i>) | 48 | 53 |
| CRUSTACEA | 22 | 1 |
| Crayfish (Cambarinae) | 11 | 1 |
| Prawns (<i>Palaemonetes kadiakensis</i>) | 11 | < 1 |
| INSECTA | 63 | 6 |
| Coleoptera | 4 | < 1 |
| Hemiptera | 19 | 1 |
| Odonata | 48 | 2 |
| Orthoptera | 26 | 3 |
| ARACHNIDA | 22 | 1 |
| Water spiders (<i>Dolomedes</i> sp.) | 22 | 1 |